

Evidence Briefing

Improving outcomes for residents of care homes

May 2016



Key messages

Providers of care in care homes should:

- offer staff training regarding recognition of cognitive states including delirium, supporting meaningful activity, end of life care
- have clear processes in place for access to healthcare
- provide information regarding dementia diagnosis and prognosis
- ensure advance care planning takes place
- offer opportunities to take part in meaningful activity

Commissioners of old peoples' services should:

- support the delivery of Comprehensive Geriatric Assessment (CGA)
- ensure broad involvement of stakeholders including care home providers and patients or representatives to ensure services are tailored
- ensure service specifications include interventions such as cognitive reframing
- ensure services are available and coordinated in relation to end of life care

- ensure stable leadership and clear strategy of overall system
- use of specific models of care can have benefit
- do not neglect long term population strategies to support healthy ageing and prevent increasing disease burden
- support the development of relationships between care homes and service providers, and ensure clear lines of responsibility
- commission targeted training for staff
- ensure evaluation and monitoring is fed back for continuous improvement

Background

Approximately 400,000 people live in care homes in England, a large proportion of whom have restricted mobility, dementia, incontinence and other medical issues¹. There are three times as many beds in care homes (both residential and nursing homes) as there are in National Health Service (NHS) hospitals.

How care homes work with health and social care services (and vice versa) is an important indicator of system integration. Historically, integration has been poor with marginalisation from NHS services^{2,3,4,5}, particularly those that offer specialist expertise in dementia,

rehabilitation and end of life care. The Atlas of Variation reports a 604-fold variation between CCGs in admissions to hospital for care home residents⁶. This issue of Evidence Briefing provides an overview of the evidence around optimising care to care home residents - to improve their health and well-being and to reduce the need for unscheduled care.

Comprehensive Geriatric Assessment

There is robust evidence that Comprehensive Geriatric Assessment (CGA) improves outcomes for vulnerable older people⁷. CGA involves a multidimensional and multidisciplinary approach which generates a problem list designed to tackle the multifactorial, interdependent issues. It establishes goal-orientated management.

Whole system care

Evidence suggests that relationships between care homes and healthcare providers are important in delivering high quality care^{2,3}. Continuity of roles and clear lines of responsibility, as well as allocated time for collaboration could help. Specialist input is important such as in undertaking a CGA. Governance and contractual arrangements have been highlighted as being of importance. Clearly defined processes could ensure roles and responsibilities are made explicit leading to improved coordination. Barriers may include high staff turnover, previous experiences and a lack of capacity.

Delivery models

A 'virtual ward' or 'hospital at home' provides an integrated health and social care team a method of delivery for CGA in the community. There is evidence for improved patient satisfaction and reduction in emergency care requests but not for reductions in admissions to hospital overall⁸.

Another approach is multi-disciplinary teams

(MDTs) with case management. A review reported four stages to this approach: identifying patients, assessing their needs, planning the care they require and responding to rapid changes⁹. Clear protocols are also required to facilitate implementation of care pathways. There is some evidence for improved outcomes for care home residents and reduced hospital admissions but the impact on overall costs is not clear.

Telemedicine involves the use of video technology to facilitate a face to face consultation without the need for patients to travel. There is some evidence to support the use of telemedicine for heart failure and diabetes etc. but no evidence specifically for care home residents^{10,11}.

Staff education

There is no conclusive research to support a particular nursing model or skill-mix¹². Poor staff knowledge is associated with increased GP referrals. Three educational priorities that have been identified are: dementia care, personal care and long term conditions¹³.

Improving use of medications

The multiple co-morbidities in this patient group result in polypharmacy (>4 drugs) which increases the likelihood of drug interactions and adverse effects. Approximately 7 out of 10 care home residents experience at least one drug error partly due to polypharmacy but also poor documentation¹⁴. A review of studies involving care home residents found no evidence that medication review, case conferencing, education or technology to support decisions improved mortality, adverse drug events or admission to hospital¹⁵. NICE guidance on this topic makes recommendations on policies, monitoring of incidents, updating and sharing information, self-administration when possible and staff training¹⁶.

Preventing pressure ulcers

Care home residents are particularly at risk of developing pressure ulcers (high prevalence of cognitive impairment and/or impaired mobility). Moore et al¹⁷ concluded that there was no reliable evidence that using risk assessment tools reduced the risk of the incidence of pressure ulcers. NICE guidance on pressure ulcer prevention and management recommended that multicomponent care plans implemented following a risk assessment are most effective at reducing pressure ulcers¹⁸. These could include pressure-relieving surfaces, repositioning of patient, skin inspections, continence management and nutrition.

Falls prevention

Falls are common in care home residents. A Cochrane Review on multi-component falls prevention programmes for care home residents suggests a possible reduction in the fall rate¹⁹. In the care home population vitamin D supplementation is effective in reducing the rate of falls but there is inconsistent evidence regarding exercise programmes. Hip protectors may reduce the risk of hip fractures in older people in residential care with a slight increased risk of pelvic fractures²⁰. Acceptability to older people and concordance are barriers to use.

Cognition, mental & physical wellbeing

NICE has published quality standards to improve the mental wellbeing (optimum independence, feeling in control, optimism, having a purpose and sense of belonging) for care home residents²¹. These standards emphasise the need for meaningful activity, the maintenance of personal identity, the recognition of possible mental illness, addressing needs related to sensory impairment, the ongoing monitoring of physical symptoms and access to healthcare. The College of Occupational Therapists has

a toolkit to support the implementation of meaningful activity in care homes, including suggestions for activities such as music, movement or group activities²². A Cochrane review reported that physical rehabilitation for care home residents may reduce disability with few adverse events, but the effects were small and may not be applicable to all residents²³.

Advance Care Planning

Advance Care Planning (ACP) is of particular relevance in care homes as half of care home residents die within approximately 1.5 years after admission²⁴. Advance care planning has been shown to result in a lower likelihood of dying in hospital and improved communication, alongside potential cost savings²⁵.

Improving care at the end of life

The NICE End of Life Care guideline applies to all care settings²⁶. It highlights the need for training in the diagnosis of dying and end of life care, to have a senior clinician responsible for the patient making decisions in conjunction with patients and relatives, to ensure adequate hydration, to distinguish between discussions around Cardiopulmonary Resuscitation and the implementation of end of life pathways, and to ensure equipment and staff are well coordinated, including out of hours.

The Gold Standards Framework is a set of standards designed to ensure high quality end of life care in primary care in use in the UK since 2001²⁷. A critical review suggests the framework is effective at improving coordination and communication, and recording patient preference regarding place of death – although evidence was limited, and use of the framework displays significant variation across practices. Ensuring GP engagement, availability, and a champion to drive the implementation of the standards were highlighted as important measures. Qualitative evidence also suggests that community matrons and district nursing

teams can play a vital part in the use of the framework with complementary strengths and expertise though a lack of clarity regarding roles and poor relationships could have a negative impact²⁸.

Last days of life

The NICE End of Life Care guideline makes specific recommendations on care in the last few days of life. It emphasises the identification of dying, communication, shared decision making, hydration, pharmacological symptom control and anticipatory prescribing²⁹.

References

1. British Geriatrics Society Joint Working Party; Quest for Quality: An Inquiry into the Quality of Healthcare Support for Older People in Care Homes: A Call for Leadership, Partnership and Improvement. BGS, 2011
2. Goodman C. Care homes and health services: an uneasy alliance. *Journal of Health Services Research & Policy*. 2016, Vol. 21(1) 1–3
3. Gage H, Dickinson A, Victor C, et al. Integrated working between residential care homes and primary care: a survey of care homes in England. *BMC Geriatr* 2012; 12: 71.
4. Glendinning C, Jacobs S, Alborz A, et al. A survey of access to medical services in nursing and residential homes in England. *Br J Gen Pract* 2002; 52: 545–548.
5. Iliffe S, Davies SL, Gordon AL, et al. Provision of NHS generalist and specialist services to care homes in England: review of surveys. *Prim Health Care Res Dev* 2015; 5 : 1 –1 6.
6. NHS Atlas of variation; NHS England, Public Health England and NHS Right Care, 2015
7. Welsh, T J, A L Gordon, and J R Gladman; Comprehensive Geriatric Assessment – a Guide for the Non-Specialist. *International Journal of Clinical Practice* 68.3 290–293, 2014
8. Shepperd S, Doll H, Angus RM, Clarke MJ, Iliffe S, Kalra L, Ricauda NA, Wilson AD. Hospital at home admission avoidance. *Cochrane Database of Systematic Reviews* 2008, Issue 4. Art. No.: CD007491. DOI: 10.1002/14651858.CD007491
9. Hong et al. Caring for High-Need, High-Cost Patients: What Makes for a Successful Care Management Program? The Commonwealth Fund, 2014.
10. Steventon et al. Effect of telehealth on use of secondary care and mortality: findings from the Whole System Demonstrator cluster randomised trial.: *BMJ* 2012;344:e3874
11. Flodgren G, Rachas A, Farmer AJ, Inzitari M, Shepperd S. Interactive telemedicine: effects on professional practice and health care outcomes. *Cochrane Database of Systematic Reviews* 2015, Issue 9. Art. No.: CD002098. DOI:10.1002/14651858.CD002098.pub2
12. Hodgkinson B, Haesler EJ, Nay R, O'Donnell MH, McAuliffe LP. Effectiveness of staffing models in residential, subacute, extended aged care settings on patient and staff outcomes. *Cochrane Database of Systematic Reviews* 2011, Issue 6. Art. No.: CD006563. DOI: 10.1002/14651858.CD006563.pub2
13. Spilsbury et al. Supporting nursing in care homes. RCN Foundation. 2015
14. Barber et al. Care homes' use of medicines study: prevalence, causes and potential harm of medication errors in care homes for older people *Qual Saf Health Care* 2009;18:341-346 doi:10.1136/qshc.2009.034231, 2008
15. Purdy. Interventions to reduce unplanned hospital admission: a series of systematic reviews. University of Bristol, 2012
16. NICE [SC1] Managing medicines in care homes. March 2014.
17. Moore ZEH, Cowman S. Risk assessment tools for the prevention of pressure ulcers. *Cochrane Database of Systematic Reviews* 2014, Issue 2. Art. No.: CD006471. DOI: 10.1002/14651858.CD006471.pub3
18. NICE CG179 Pressure ulcers: prevention and management. NICE, 2016
19. Cameron ID, Gillespie LD, Robertson MC, Murray GR, Hill KD, Cumming RG, Kerse N. Interventions for preventing falls in older people in care facilities and hospitals. *Cochrane Database of Systematic Reviews* 2012, Issue 12. Art. No.: CD005465. DOI: 10.1002/14651858.CD005465.pub3
20. Santesso N, Carrasco-Labra A, Brignardello-Petersen R. Hip protectors for preventing hip fractures in older people. *Cochrane Database of Systematic Reviews* 2014, Issue 3. Art. No.: CD001255. DOI: 10.1002/14651858.CD001255.pub5
21. NICE QS50, Mental wellbeing of older people in care homes, 2013
22. College of Occupational Therapists. 201 5. Living well through activity in care homes: The toolkit. <https://www.cot.co.uk/sites/default/files/general/public/Unit2-Care-home-staff-resources-2015.pdf>
23. Crocker T, Forster A, Young J, Brown L, Ozer S, Smith J, Green J, Hardy J, Burns E, Glidewell E, Greenwood DC. Physical rehabilitation for older people in long-term care. *Cochrane Database of Systematic Reviews* 2013, Issue 2. Art. No.: CD004294. DOI:10.1002/14651858.CD004294.pub3
24. Forder, J. and Fernandez, J.L. Length of stay in care homes. PSSRU Discussion Paper 2769, 2011
25. Halpern, S. (2015) Towards evidence-based end-of-life care. *N Engl J Med* 2015; 373:2001-2003
26. NICE QS13 End of life care for adults, 2011
27. Shaw, K.L. et al. Improving end-of-life care: a critical review of the Gold Standards Framework in primary care. *Palliative Medicine*, 2010
28. King, N. et al. Community nursing roles and the Gold Standards Framework for community palliative care. Centre for Applied Psychological Research, School of Human and Health Sciences, University of Huddersfield, 2008
29. NG31 Care of dying adults in the last days of life, 2015